

IOWA STATE UNIVERSITY

Digital Repository

Plant Pathology and Microbiology Publications

Plant Pathology and Microbiology

8-9-2019

A new mechanism for translational control in plants

W. Allen Miller

Iowa State University, wamiller@iastate.edu

Savithramma P. Dinesh-Kumar

University of California, Davis

Follow this and additional works at: https://lib.dr.iastate.edu/plantpath_pubs



Part of the [Genetics Commons](#), [Plant Biology Commons](#), and the [Plant Pathology Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/plantpath_pubs/266. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Response or Comment is brought to you for free and open access by the Plant Pathology and Microbiology at Iowa State University Digital Repository. It has been accepted for inclusion in Plant Pathology and Microbiology Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

A new mechanism for translational control in plants

Abstract

In unstressed conditions, the cap-binding protein, eIF4E, binds the 5' cap structure (m7G) on mRNA and recruits other translation initiation factors to bring the ribosome to the mRNA. Bruns *et al.* show that, in plants, phosphorylation of eIF4E by the master metabolic regulatory protein, sucrose nonfermenting-related kinase 1 (SnRK1), reduces translation globally. This negative regulation of translation inhibits geminivirus infection and is speculated to be a response to various abiotic stresses.

Disciplines

Genetics | Plant Biology | Plant Pathology

Comments

This is the peer reviewed version of the following article: Miller, W. Allen, and Savithramma P. Dinesh-Kumar. "A new mechanism for translational control in plants." *The FEBS Journal*, which has been published in final form at doi: [10.1111/febs.15022](https://doi.org/10.1111/febs.15022). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Received Date : 03-Jul-2019
Accepted Date : 27-Jul-2019
Article type : Commentary

Color : Fig 1

Commentary: FEBS Journal

A new mechanism for translational control in plants

W. Allen Miller¹ and Savithramma P. Dinesh-Kumar²

¹Department of Plant Pathology & Microbiology, Iowa State University, Ames, IA, USA

²Department of Plant Biology and The Genome Center, College of Biological Sciences, University of California Davis, Davis, CA, USA

Correspondence:

W. Allen Miller, Department of Plant Pathology & Microbiology, Iowa State University, 2213 Pammel Drive, Ames, IA 50011, USA
E-mail: wamiller@iastate.edu

Abbreviations

eIF4E, eukaryotic initiation factor 4E
eIFiso4E, eukaryotic initiation factor iso4E
4E-BP, eIF4E-binding protein
PKR, protein kinase R
SNF1, sucrose non-fermenting 1
SnRK1, SNF1-related kinase 1
AMPK, AMP-activated kinase
SnRK1-KD, SnRK1 kinase domain
SnRK1-KDKR, SnRK1 kinase domain inactive
BiFC, bimolecular fluorescent complementation

The process of protein synthesis is well-conserved among eukaryotes [1]. However, regulation of translation differs between plants and animals [2]. One means of translational control

involves eukaryotic initiation factor 4E (eIF4E), which binds the 5' cap structure of mRNA, and its partner eIF4G to recruit other initiation factors and ultimately the ribosome to the mRNA. Plants also contain an isoform, eIFiso4E, which pairs with eIFiso4G. The different biological roles of eIF4E/4G vs eIFiso4E/iso4G are unclear, but *Arabidopsis thaliana* (At) plants lacking eIFiso4E or eIFiso4G show developmental abnormalities [3]. In mammals, under low nutrient or various stress conditions, eIF4E-binding protein (4E-BP) inhibits translation initiation by preventing eIF4E from binding eIF4G and thus from recruiting the other factors to the mRNA (Figure 1) [4]. Excessive translation initiation due to aberrant lack of inhibition of eIF4E by 4E-BP can lead to many types of cancer and metabolic disorders [5]. While in plants and animals eIF4E is structurally similar [6], plants lack an ortholog to the mammalian 4E-BP [7]. However, Patrick et al. [8] identified a different eIF4E binding protein (CBE1) in *Arabidopsis* that negatively regulates accumulation of mRNAs that control the cell cycle and mitosis, via a mechanism that is currently still unknown.

In vertebrates, translation is greatly attenuated during virus infection. This inhibits virus gene expression because viruses depend on the host's translation machinery. During virus infection, protein kinase R (PKR) becomes activated by binding to double-stranded RNA, which is generated during RNA virus replication, and it then phosphorylates eIF2 α [9]. This prevents the exchange of GDP for GTP required for regeneration of the ternary complex eIF2•GTP•tRNA^{Met}, necessary for initiation of translation (Figure 1). Plants lack a PKR homolog, and do not appear to use this mechanism as an antiviral response [10]. Given that translation attenuation is important for rapid response to changes in the environment or during virus infection, it is likely that such regulation takes place in plants by mechanisms that have yet to be discovered.

Indeed, a new mechanism of translation attenuation, apparently unique to plants, has been reported by the Bisaro lab in this issue of *The FEBS Journal* [11]. They provide evidence for translation attenuation mediated by phosphorylation of eIF4E and eIFiso4E by sucrose non-fermenting (SNF)-related kinase 1 (SnRK1). SnRK1 is an ortholog of yeast SNF1 and mammalian AMP-activated kinase (AMPK) that functions as a master regulator of energy homeostasis [12]. During biotic or abiotic stress, SNF1/AMPK1/SnRK1 activates energy-generating catabolic pathways and inhibits energy-consuming anabolic pathways, at the transcriptional level and post-translationally, by phosphorylation of key enzymes [13]. Here, Bruns et al identified two consensus SnRK1 target sites in *Arabidopsis* eIF4E and eIFiso4E that are absent in mammalian eIF4E. The purified SnRK1 kinase domain (SnRK1-KD), but not a catalytically inactive mutant (SnRK1-KDKR), phosphorylates the predicted target sites in At eIFiso4E. Consistent with this, SnRK1 interacts with At eIF4E and At eIFiso4E in a yeast two-hybrid assay and in bimolecular fluorescent complementation (BiFC) assays in plant cells. Furthermore, SnRK1 phosphorylates At eIFiso4E when both are transiently co-expressed in plant cells.

To determine the effect of SnRK1-mediated phosphorylation on At eIF4E/iso4E activity, the authors expressed SnRK1 under a copper inducible promoter in a yeast strain that lacks endogenous eIF4E (*cdc33Δ*) but expresses At eIF4E, At eIFiso4E or human eIF4E (*Hs* eIF4E). Interestingly, induction of SnRK1 increased doubling time of yeast that depended on At eIF4E or At eIFiso4E but not *Hs* eIF4E. This reduced growth of yeast is likely due to phosphorylation of At

eIF4E/iso4E by SnRK1 because *Hs* eIF4E lacks the SnRK1 phosphorylation site present on plant eIF4E/iso4E. In the SnRK1-growth-inhibited yeast, there was a substantial reduction in translation as measured by a 50% reduction in the ratio of polysomes (ribosome-studded, actively translating mRNAs) to non-polysomes (non-translating ribosomes). However, as predicted, no such SnRK1-mediated reduction in translation was observed in yeast complemented with *Hs* eIF4E or non-phosphorylatable mutant *At* eIF4E-T67V/T91A (called *At* eIF4E-VA). A similar reduction in polysome/non-polysome ratio was also observed in *Nicotiana benthamiana* plants engineered to overexpress SnRK1. Finally, transient overexpression of *At* eIF4E or *At* eIF4E-VA in SnRK1-overexpressing plants restored the polysome/non-polysome ratio to that found in wild type plants, presumably because the amount of overexpressed eIF4E was sufficient to allow full translation. As expected, overexpression of the phosphomimic mutant, *At* eIF4E-T67D, had no effect on the reduced polysome/non-polysome ratio, because it behaves as a phosphorylated (inactivated) eIF4E under all conditions.

The results described in Bruns et al. [11] provide compelling evidence for translation attenuation due to SnRK1-mediated phosphorylation of eIF4E and eIFiso4E. However, many questions remain. How does phosphorylation inhibit eIF4E/iso4E activity (Figure 1)? Does it affect binding to eIF4G or to the 5' cap on the mRNA or other components of the translation machinery? Does SnRK1-mediated phosphorylation of eIF4E functionally replace the role that 4E-BP plays in mammals? If it does, then the phosphorylation state of eIF4E should increase under nutrient stress/low energy conditions in plants. Does SnRK1-mediated translation inhibition play an antiviral role? Previously, Bisaro's group showed that SnRK1 overexpression increases resistance to a geminivirus, while SnRK1 knockdown makes plants more susceptible to the virus [14]. Therefore, translation attenuation would inhibit geminivirus gene expression because its mRNAs, generated by host transcription machinery from the viral DNA, are translated by normal cellular mechanisms, unlike many RNA viruses that are translated cap-independently [15] and thus may not be subject to eIF4E regulation. How is translation of individual mRNAs affected by eIF4E phosphorylation? It's unlikely that all are affected equally. One would predict that mRNAs needed during low energy conditions would translate more efficiently than those that are not. Ribosome profiling [16] could be one approach to answer this question.

In summary, Bruns et al. have provided a first answer to the complex question of how plants negatively control translation, identifying a novel regulatory mechanism that is unknown in mammals. How SnRK1-mediated eIF4E phosphorylation is regulated, and the biological pathways controlled by this mechanism remain to be determined.

Acknowledgement. The authors thank Karen Browning for advice. Research on virus-eIF4E interactions proteins in WA Miller's laboratory has been supported by the US National Institutes of Health (R01 GM067104). Research on host-virus interactions in Dinesh-Kumar laboratory is supported by US National Science Foundation (IOS-1339185) and National Institute of Health (R01 GM132582)

Conflict of interest statement. The authors declare no conflicts of interest.

References

1. Jackson, R. J., Hellen, C. U. & Pestova, T. V. (2010) The mechanism of eukaryotic translation initiation and principles of its regulation, *Nat Rev Mol Cell Biol.* **11**, 113-127.
2. Browning, K. S. & Bailey-Serres, J. (2015) Mechanism of cytoplasmic mRNA translation, *The Arabidopsis book / American Society of Plant Biologists.* **13**, e0176.
3. Lellis, A. D., Allen, M. L., Aertker, A. W., Tran, J. K., Hillis, D. M., Harbin, C. R., Caldwell, C., Gallie, D. R. & Browning, K. S. (2010) Deletion of the eIFiso4G subunit of the Arabidopsis eIFiso4F translation initiation complex impairs health and viability, *Plant Mol Biol.* **74**, 249-63.
4. Richter, J. D. & Sonenberg, N. (2005) Regulation of cap-dependent translation by eIF4E inhibitory proteins, *Nature.* **433**, 477-80.
5. Musa, J., Orth, M. F., Dallmayer, M., Baldauf, M., Pardo, C., Rotblat, B., Kirchner, T., Leprivier, G. & Grunewald, T. G. (2016) Eukaryotic initiation factor 4E-binding protein 1 (4E-BP1): a master regulator of mRNA translation involved in tumorigenesis, *Oncogene.* **35**, 4675-88.
6. Monzingo, A. F., Dhaliwal, S., Dutt-Chaudhuri, A., Lyon, A., Sadow, J. H., Hoffman, D. W., Robertus, J. D. & Browning, K. S. (2007) The structure of eukaryotic translation initiation factor-4E from wheat reveals a novel disulfide bond, *Plant Physiol.* **143**, 1504-1518.
7. Browning, K. S. (2004) Plant translation initiation factors: it is not easy to be green, *Biochem Soc Trans.* **32**, 589-591.
8. Patrick, R. M., Lee, J. C. H., Teetsel, J. R. J., Yang, S. H., Choy, G. S. & Browning, K. S. (2018) Discovery and characterization of conserved binding of eIF4E 1 (CBE1), a eukaryotic translation initiation factor 4E-binding plant protein, *J Biol Chem.* **293**, 17240-17247.
9. Bou-Nader, C., Gordon, J. M., Henderson, F. E. & Zhang, J. (2019) The search for a PKR code-differential regulation of protein kinase R activity by diverse RNA and protein regulators, *RNA.* **25**, 539-556.
10. Bilgin, D. D., Liu, Y., Schiff, M. & Dinesh-Kumar, S. P. (2003) P58(IPK), a plant ortholog of double-stranded RNA-dependent protein kinase PKR inhibitor, functions in viral pathogenesis, *Dev Cell.* **4**, 651-661.
11. Bruns, A. N., Li, S., Mohannath, G. & Bisaro, D. M. (2019) Phosphorylation of Arabidopsis eIF4E and eIFiso4E by SnRK1 inhibits translation, *Febs j.*
12. Emanuelle, S., Doblin, M. S., Stapleton, D. I., Bacic, A. & Gooley, P. R. (2016) Molecular Insights into the Enigmatic Metabolic Regulator, SnRK1, *Trends Plant Sci.* **21**, 341-353.
13. Hulsmans, S., Rodriguez, M., De Coninck, B. & Rolland, F. (2016) The SnRK1 Energy Sensor in Plant Biotic Interactions, *Trends Plant Sci.* **21**, 648-661.
14. Hao, L., Wang, H., Sunter, G. & Bisaro, D. M. (2003) Geminivirus AL2 and L2 proteins interact with and inactivate SNF1 kinase, *Plant Cell.* **15**, 1034-48.
15. Miras, M., Miller, W. A., Truniger, V. & Aranda, M. A. (2017) Non-canonical Translation in Plant RNA Viruses, *Front. Plant Sci.* **8**, 494.
16. Hsu, P. Y., Calviello, L., Wu, H. L., Li, F. W., Rothfels, C. J., Ohler, U. & Benfey, P. N. (2016) Super-resolution ribosome profiling reveals unannotated translation events in Arabidopsis, *Proc Natl Acad Sci U S A.*

Figure Legend

Figure 1. Negative regulation of translation in mammals and plants.

Top: In normal, unstressed conditions, cap-binding protein eIF4E simultaneously binds the m⁷GpppN cap structure on the mRNA and scaffolding protein eIF4G, which binds other initiation factors (OIFs) to recruit the 43S complex consisting of the 40S ribosomal subunit bound to the ternary complex of eIF2•GTP•tRNA_i^{Met}. *Middle:* Under low nutrient or stress conditions, eIF4E binding protein (4E-BP) binds eIF4E to prevent eIF4G binding, thus blocking translation initiation. In response to many virus infections, PKR phosphorylates eIF2 α , preventing exchange of GTP with GDP in eIF2B, which, in turn, prevents formation of the ternary complex. *Bottom:* In plants, Bruns et al provide evidence that phosphorylation of eIF4E by SnRK1 attenuates translation initiation, but the interactions disrupted by this modification are unknown (question marks).

Legend for Graphical Abstract.

In unstressed conditions, the cap-binding protein, eIF4E, binds the 5' cap structure (m7G) on mRNA and recruits other translation initiation factors to bring the ribosome to the mRNA. Bruns et al show that, in plants, phosphorylation of eIF4E by the master metabolic regulatory protein, sucrose non-fermenting-related kinase 1 (SnRK1), reduces translation globally. This negative regulation of translation inhibits geminivirus infection and is speculated to be a response to various abiotic stresses.

Type of file: figure

Label: Figure 01

Filename: febs_15022_f1.pdf





Figure 1

Type of file: figure

Label: Graphic Abstract Fig

Filename: febs_15022_G_Abstract_Fig.pdf

Normal conditions



Geminivirus infection / Abiotic stresses?

